

Effects of Glibenclamide and Metformin on serum uric acid levels in patients with type 2 diabetes mellitus

Najlaa Saadi Ismail

Department of Pharmacology, Mosul College of Medicine, University of Mosul

Received Accepted

13.7.2010 30.11.2010

ABSTRACT

Objectives: To assess the effect of Glibenclamide and Metformin on serum uric acid level in patient with type 2 diabetes.

Study design: case control study.

Study period: from March 2009 to January 2010.

Subjects and Methods: Fasting blood sugar and serum uric acid level were measured in patient suffering from type 2 diabetes mellitus who were referred to Al-Wafa Diabetic Center in Mosul City: (group 1) 32 patients on glibenclamide therapy, (group 2) 42 patients on metformin therapy and (group 3) 42 patients on combination therapy, (group 4) 32 patients on restricted diet, and 23 apparently healthy volunteers, were taken as a control group.

Results: Study showed a significant increase in the serum uric acid level of the diabetic patients as compared with the control. glibenclamide and/or metformin showed no significant difference in the serum uric acid level in patients with type 2 diabetes mellitus.

Conclusion: glibenclamide and / or metformin had no significant effect on serum uric acid level in patients with type 2 diabetes mellitus.

Key Words: Uric Acid, Hyperuricemia, type 2 diabetes mellitus, glibenclamide, metformin.

الخلاصة

أهداف البحث: لدراسة تأثير عقار الميتغورمين و عقار الكلابينكلاميد على حامض البيريك في مصل المرضى المصابين بداء السكري النمط الثاني.

تصميم الدراسة: دراسة عينية مقارنة.

فترة الدراسة: من آذار ٢٠٠٩ إلى كانون الثاني ٢٠١٠.

الطرق المتبعة والأشخاص: تم قياس تركيز الكلوكوز في مصل الدم وقياس مستوى حامض البيريك في مصل الدم لمجموعات المرضى المصابين بداء السكري النمط الثاني اللذين يراجعون مركز الوفاء للسكري :

(المجموعة الأولى) ٣٢ مريض تحت علاج الكلابينكلاميد (المجموعة الثانية) مريض ا

ميتغورمين، (المجموعة الثالثة) مريض تحت علاج الكلابينكلاميد والميتغورمين معا (

) مريض على الحمية و مجموعة الضبط مريض من الاصحاء غير السكريين .

النتائج: اظهر مرض السكري النمط الثاني ارتفاعا في حامض البيريك كمقارنة مع مجموعة الضبط لم يظهر عقار الميتغورمين او الكلابينكلاميد أي تأثير معنوي على مستوى حامض البيريك في مصل المرضى المصابين بداء السكري النمط الثاني .

الاستنتاج: يوجد تأثير للميتغورمين او الكلابينكلاميد على مستوى حامض البيريك في مصل الدم في مجموعات المرضى المصابين بداء السكري النمط الثاني .

Uric acid is the final catabolic, heterocyclic purine derivative resulting from the oxidation of purines in humans. Due to the loss of hepatic

uricase activity during human evolution, uric acid is excreted as such and is not further metabolized into carbon dioxide and ammonia. A major

mechanism underlying hyperuricemia is impaired renal excretion of urate¹. There has been growing interest in the association of hyperuricemia with hyperglycemia². uric acid may be a marker of oxidativestress³ and may have a potential therapeutic role as an antioxidant^{4,5}. Like other strong reducing substances such as ascorbate, uric acid can also act as a prooxidant particularly at elevated levels.

Patients with non-insulin-dependent diabetes mellitus (NIDDM) are at increased risk for cardiovascular diseases such as hypertension and stroke. Hyperuricemia is a common finding in NIDDM, but its significance as an independent risk factor for cardiovascular disease has remained uncertain^{6,7}, serum uric acid level is positively associated with the development of type 2 diabetes regardless of various study characteristics. Further research should attempt to determine whether it is effective to utilize serum uric acid level as a predictor of type 2 diabetes for its primary prevention⁸.

The association of high serum uric acid with insulin resistance has been known since the early part of the 20th century, nevertheless, recognition of high serum uric acid as a risk factor for diabetes has been a matter of debate. In fact, Hyperuricemia has always been presumed to be a consequence of insulin resistance rather than its precursor⁸. Elevated levels of uric acid should alert physicians to the possibility of insulin resistance. The serum uric acid level was associated with insulin resistance and plasma glucose levels more strongly in females than in males in our study population⁹. serum uric acid

is positively associated with serum glucose in healthy subjects¹⁰.

Evidence has accumulated indicating that the generation of reactive oxygen species (oxidative stress) may play an important role in the etiology of diabetic complications. This hypothesis is supported by evidence that many biochemical pathways strictly associated with hyperglycemia (glucose autoxidation, polyol pathway, prostanoid synthesis, protein glycation) can increase the production of free radicals¹¹.

Drugs with other primary uses, that have known uricosuric properties which decrease serum uric acid levels, such as losartan¹², fenofibrate^{13,14}, whereas diuretic¹⁵ pyrazinamide¹⁶, elevate serum uric acid levels.

The present study was conducted to investigate the effect of glibenclamide and metformin on serum uric acid level in a number of type 2 diabetic patients.

Patients and methods

This is a case-control study which was conducted in the Department of Pharmacology, College of Medicine, Univrsity of mosul and Al-Wafa Diabetic Center in Mosul from March, 2009 to January, 2010. Five groups were enrolled for this study: the first group included 32 type 2 diabetic patients, their ages ranged between (32-70 years) treded with glibenclamide, the second group included 42 type 2 diabetic patients, their ages ranged between (27-75 years) treated with metformin, the third group included 42 type 2 diabetic patients, their ages ranged between (42-66 years) treated with combination of these two drugs (glibenclamide and metformin), the fourth group included

32 type 2 diabetic patients on restricted diet therapy only, their ages ranged between (28-85 years) and the fifth group included 23 apparently healthy volunteers participated as a control group, their ages ranged between (28-56 years).

Five ml of venous blood samples were collected from each individual (patient and control) after at least 12 hours fasting. Fasting blood sugar was measured using a glucose oxidase method¹⁷ which is available as a kit manufactured by Biomaghreb. Serum uric acid was assessed by uricase enzymatic method¹⁸ by using a special kit (Biolabo).

Statistical analysis

The values were quoted as mean \pm SE. P-values <0.05 was considered to be statistically significant. Unpaired- test was used to compare the data obtained in this study.

Results

Table 1 shows the characteristics of the study participants. The table shows the ages of the individual and the number of males and females in the five groups.

Comparison between fasting blood sugar and serum uric acid level of Glibenclamide, Metformin, combination and diet groups with those of the control showed a significant elevation of all parameters. Comparison of fasting blood sugar and serum uric acid level of glibenclamide, metformin and combination groups with those of diet showed a significant elevation for fasting blood sugar and a non significant differences for serum uric acid level. Comparison between fasting blood sugar and serum uric acid level of glibenclamide and metformin, no significant differences were found between them. (Table 2)

Table 1. The characteristic of study participants

CHARACTERISTIC	Control group N=23	Glibenclamide group N=32	Metformin group N=42	Combination group N=42	Diet group N=32
Sex M	8	21	20	23	19
F	15	11	22	19	13
Age (year)	28-56 45 ± 2.4	32-70. 50 ± 2.0	27-75 47 ± 1.6	42-66 44 ± 1.3	28-85 50 ± 2.1

Table 2. The comparison between different groups

Parameters	Control group	Glibenclamide group	Metformin group	Combination group	Diet group
Fasting blood sugar (mmol/l) All groups and Control	5.32±0.24	12.87±1.28 **	11.02±0.75 **	10.97±0.62 **	9.74±0.74 **
Serum uric acid (μmol/l) All groups and Control	265.95±17.52	404.18±27.39 **	470±61.01 *	452±38.35 **	461±28.39 **
Fasting blood sugar (mmol/l) All groups and Diet	---	12.87±1.28 *	11.02±0.75 **	10.97±0.62 *	9.74±0.74
Serum uric acid (mmol/l) All groups and Diet	---	404.18±27.39 P=0.15 (NS)	470.04±61.01 P=0.9 (NS)	452±38.35 P=0.85 (NS)	461±28.39
Fasting blood sugar (mmol/l) Glibenclamide and Metformin	---	12.87±1.28 P=0.219 (NS)	11.02±0.75 P=0.219 (NS)	---	---
Serum uric acid (mmol/l) Glibenclamide and Metformin	---	404.18±27.39 P=0.32 (NS)	470.04±61.01 P=0.32 (NS)	---	---

(NS) Non significant, * P <0.05, ** P <0.01

Discussion

Hyperuricemia has previously been described as strong predictor of well defined cerebrovascular complications in patients with type 2 diabetes⁷.

The present study showed a significant elevation of serum uric acid level of the diabetic patients as compared with the control individuals. The patients are hyperglycemic, as evident by the high concentration of fasting blood sugar.

Variations in uric acid levels have been increasingly associated with

insulin resistance, hyperinsulinemia, and diabetes^{19, 20}. In 2009, a study was conducted by kodama and saito⁸ they reported that uric acid levels are higher in subjects with prediabetes and early type 2 diabetes than in healthy. In type 2 diabetes, Hyperuricemia seems to be associated with the insulin-resistant syndrome and with early onset or increased progression to overt nephropathy, while hypouricemia is associated with worse metabolic control, hyperfiltration and a late onset or decreased progression to overt nephropathy²¹, Interestingly, serum

uric acid levels were increased in type 2 diabetic patients and this phenomenon seemed to be more profound in male diabetic patients²².

This study is in contrast to the study conducted by Gotoh et al, they reported that serum uric acid levels in diabetics are significantly lower than those in non-diabetic subjects²³.

The finding of this study showed no significant difference between serum uric acid levels of diet group with glibenclamide, metformin group, this findings are similar to those of Luque et al, where they reported that there was no change in serum uric level observed with metformin for treatment of polycystic ovary syndrome²⁴, so as the study done by Fruehwald and Oltmanns²⁵.

In contrast to these results, Gregorio and Manfrini in 1996, reported that metformin lowered uric acid in elderly type 2 diabetic patients²⁶. Gokcel and Gumurdulu in research concerning the evaluation of the safety and efficacy of metformin in the treatment of obesity, they reported that metformin administration resulted in a significant reduction in serum uric acid levels in obese patients²⁷. Barskova et al reported that metformin reduce production of uric acid in patient with gout and insulin resistant²⁸.

The present study revealed that glibenclamide had no significant effect on serum uric acid levels in patients with type 2 diabetes. These finding are similar to the studies conducted by Cheach²⁹, and Carvalho³⁰. Their findings might indicate that, hypoglycemic agents have no effect on the serum uric acid level of the diabetic patients and does not affect the balance

between urate production and renal excretion.

Conclusion glibenclamide, metformin or their combination had no significant effect on serum uric acid level in patient with type 2 diabetes mellitus as evident by non significant differences between patients taking these drugs and patients on diet only.

References

1. Melanie K, Serena S. Meta-Analysis of 28,141 Individuals Identifies Common Variants within Five New Loci That Influence Uric Acid Concentrations. *PLoS Genet.* 2009; 5(6): e1000504.
2. Nan H, Pang Z, Wang S .Serum uric acid, plasma glucose and diabetes. *Diab Vasc Dis Res* 2010; 7(1):40-46.
3. Becker BF. Towards the physiological function of uric acid. *Free Radical Biol Med* 1993; 14 (6): 615-631.
4. Glantzounis GK, Tsimoyiannis EC, Kappas AM. Uric acid and oxidative stress . *Curr Pharm Des* 2005; 11 (32): 4145–4151.
5. Bruce N., Richard C, Elizabeth S. Uric acid provides an antioxidant defense in humans against oxidant- and radical-caused aging and cancer. *Biochemistry* 1981; 78(11): 6858-6862.
6. Hyperuricemia as a risk factor on cardiovascular events in Taiwan: The Chin-Shan Community Cardiovascular Cohort Study. *Atherosclerosis* 2005; 183(1):147-155.
7. Lehto S, Niskanen L, Rönnemaa T, Laakso M .Serum uric acid is a strong predictor of stroke in patients with non-insulin-

dependent diabetes mellitus. *Stroke* 1998; 29(3):635-639.

8. Kodama S, Saito K, Yachi Y. Association between Serum Uric Acid and Development of type 2 Diabetes. *Diabetes Care* 2009; 32 (9):1737-1742.
9. Chou D, Lin H, Sia S .Gender differences in the relationships of serum uric level with fasting serum insulin and plasma glucose in patients without diabetes. *Rheumatol* 2001; 28(3):571-576.
10. Facchini F, Chen Y, Hollenbeck C. Relationship between resistance to insulin-mediated glucose uptake, urinary uric acid clearance, and plasma uric acid concentration. *Jama* 1991; 266:3008-3011.
11. Giugliano D, Ceriello A, Paolisso G, Oxidative Stress and Diabetic Vascular Complications. *Diabetes Care* 1996; 19(3): 257-267.
12. Daskalopoulou S, Tzovaras V, Mikhailidis D. Effect on Serum Uric Acid Levels of Drugs Prescribed for Indications other than Treating hyperuricaemia. *Curr Pharm Des* 2005; 11(32); 4161-4175.
13. Desager JP, Hullhoven R, Harvengt C. Uricosuric effect of fenofibrate in healthy volunteers. *J Clin Pharmacol* 1980; 20:560-564.
14. Harvengt C, Heller F, Desager JP. Hypolipidemic and hypouricemic action of fenofibrate in various types of hyperlipoproteinemias. *Artery* 1980; 7(1):73-82.
15. Reyes A. The increase in serum uric acid concentration caused by diuretics might be beneficial in heart failure. *Eur J heart failure* 2005; 7 (4):461-467.
16. Solangi G, zaberti B, shiakh S. pyrazinamide induced hyperuricemia in pateints taking anti tubercullous therapy. *JCPSP* 2004; 14 (3): 136-138.
17. Lott J, Turner K. Evaluation of trinder s glucose oxidase method for measuring glucose in serum and urine. *Clin Chem* 1975; 21:1754-1760.
18. Gochman N, Schmitz J. Automated determination of uric acid with use of an uricase-peroxidase system. *Clin Chem* 1971; 17:1154-9.1999; 34:60-63.
19. Lebto S, Niskanen L, Ronneman T, et al serum uric acid is astrong predictor of stroke in patients with type 2 diabetes mellitus .stroke:1998;29:635-639 .
20. Seghier G, Moruzzo D, Fassetti S increase in serum uric acid is selectively associated with stroke in type 2 diabetes. *Diabetes care* 2002; 25(6):1095.
21. Cavallo P, Gentile, L, Repetti, E. Hypouricemia and hyperuricemia in type 2 diabetes: two different phenotypes .*Eur J Clinical Investigation* 2001; 31(4): 318-321.
22. Casevic A, Semiz S, Macic C. Arelevance of uric acid in progression of type 2 diabetes. *bosn j basic med sci* 2010;10 (1):54-59.
23. Gotoh M, Yatoh M, Iguchi A. Serum uric acid concentrations in type 2 diabetes: its significant relationship to serum 1, 5-anhydroglucitol concentrations. *Endocr Regul* 2005; 39(4):119-25.
24. Luque M, AlvarezB, Uriol M. Serum uric acid concentration as non-classic cardiovascular risk factor in women with polycystic ovary syndrome: effect of

treatment with ethinyl-estradiol plus cyproterone acetate versus Metformin. *Hum Reproduction* 2008; 23(7):1594-601.

25. Fruehwald B, Oltmanns K, Toschek B. Short-term treatment with Metformin decreases serum leptin concentration without affecting body weight and body fat content in normal-weight healthy men. *Metabolism* 2002; 51(4):531-36.

26. Gregorio F, manfrin S. Metformin treatment in elderly type 2 diabetic patients .*Arch Gerontol Geriatr* 1996; 22(1):261-270.

27. Gokcel A, Gumurdulu Y, Karakose H. Evaluation of the safety and efficacy of sibutramine, orlistat and Metformin in the treatment of obesity. *Diabetes Obes Metab* 2002; 4 (1): 49 – 55.

28. Barskova V, Eligeov M, kadaeve F. Effect of Metformin on clinical course of gout and insulin resistant .*clin med* 2009; 87(7):41-6.

29. Cheach j. Aclinical evaluation of Glibenclamide (hb 419"daonil") anew antidiabetic sulphonylurea . *singapore med j* 1972; 13(2) ; 87-88.

30. Carvalho M, Oliveira D, Suzuki K. The action of Glibenclamide on glycogen catabolism and related parameters in the isolated perfused rat liver. *Res Commun Mol Pathol Pharmacol* 2006; 119(1):115-126 .